

Sarah Smith

### **Investigating heat loss from huddling penguins**

In my investigation, the one thing that I shall change by equal amount each time is the position of the test tube been assessed. (The position in a huddle)

In my investigation, the thing I shall measure, will be the temperature.

To make the investigation a fair test, here is a list of things that I'm going to keep the same:

- Test tubes sizes
- Amount of water in the test tubes
- Time at which we start the clock
- Starting time

**Apparatus** – here is a list of everything I shall need to do my investigation:

- Test tubes (penguins)
- Hot water
- Thermometer
- Water heater
- Stop watch
- Beakers
- Elastic band
- Goggles
- Kettle

**My prediction** – what I think will happen:

I think that the difference in position will affect the temperature. Therefore I think that the penguin positioned in the middle will have the greatest temperature all the way through the experiment.

Why I think this will happen:

A penguin will lose heat in 3 different ways. One, by conduction – where heat moves through a solid by the particles vibrating (creating the knock on effect). Two, by convection – this is where hot air/particles rise and cold air/particles rush to take its place causing convection currents. And three, by radiation. Radiation can travel through empty space and needs no form of medium.

The penguin stood on its own will lose heat faster, than the penguins in a huddle, as its temperature (heat source) is greater than the penguin's environment. The process is similar to diffusion, where there is a net movement of particles from a region of high concentration to an area of low concentration. Except, this is where heat moves more rapidly to an area where it's colder. The greater the difference between the temperatures, the faster it travels. So the heat will be transferred faster from the penguin on its own into the environment than the penguins in a huddle. However in a huddle the penguin in the middle will lose heat very slowly as its environment is the other penguins and they all virtually have the same temperature. The penguins surrounding the penguin in the middle, will lose heat, not as much as the penguin stood on its own but greater than the penguin stood in the middle. This is because some of their heat will be transferred to each other and some into the environment. Also, penguins have small feathers which they "fluff up". By doing this, they trap air which acts as an insulator – keeping heat in. So, the test tube in the middle will be insulated because of the warm air surrounding it.

**Plan** – here is what I plan to do:

First I shall set up my apparatus as shown in the diagram, and then I shall fill each test tube with 10 cm of hot water with a temperature of about 70 degrees C. I will then measure the temperature of the water, immediately, in the test tube in the middle of the huddle. At the same time, I will measure the temperature of the water in the test tubes around the huddle and the test tube standing on its own. While doing this I will make sure that everything will be kept fair, to ensure a fair test. I will record the temperature results in a results table. I will repeat the above two minutes after for 10 minutes. I will repeat the whole experiment twice to ensure correct and accurate readings. To guarantee that I have accurate readings, I will try and get the starting temperature to be the same both times. I will then make a graph from the results.

**Results**

What I changed:

What I measured: temperature Units: C (degrees)

Number of test tubes

|                  | Start     | 2mins       | 4mins       | 6mins       | 8mins       | 10mins      |
|------------------|-----------|-------------|-------------|-------------|-------------|-------------|
| Own              | 76        | 61          | 57          | 54          | 49          | 45          |
| Middle           | 76        | 67          | 64          | 63          | 61          | 60          |
| Edge             | 76        | 67          | 64          | 61          | 59          | 57          |
| Own              | 76        | 60          | 58          | 50          | 50          | 49          |
| Middle           | 76        | 71          | 69          | 67          | 61          | 59          |
| Edge             | 76        | 70          | 68          | 65          | 60          | 47          |
| Own              | 76        | 59          | 57          | 54          | 52          | 48          |
| Middle           | 76        | 73          | 69          | 67          | 65          | 62          |
| Edge             | 76        | 70          | 66          | 63          | 59          | 54          |
| <b>Average 1</b> | <b>76</b> | <b>60</b>   | <b>57.3</b> | <b>52.6</b> | <b>50.3</b> | <b>47.3</b> |
| <b>Average 2</b> | <b>76</b> | <b>70.3</b> | <b>67.3</b> | <b>65.6</b> | <b>62.3</b> | <b>60.3</b> |
| <b>Average 3</b> | <b>76</b> | <b>69</b>   | <b>66</b>   | <b>63</b>   | <b>59.3</b> | <b>52.6</b> |

### **Conclusion**

What I have found out by looking at my graph is that the penguin (test tube) standing on its own loses the heat fastest, while the penguin in the middle of the huddle loses heat more slowly. For example, the penguin on its own after 2 minutes is at the temperature 61 degrees while the penguin in the huddle is at 70 degrees. This is because the differences in the temperatures, between the penguin (test tube) and its environment was getting smaller, causing the heat to move more slowly. Also, the penguin in the middle was insulated better therefore surrounded by warm air. You can see this by the gradient of the graph decreases as the time passed showing that the rate is decreasing. Even though that the "edge position of the huddle" reading of the graph shows the temperature increasing slightly (gaining heat which obviously isn't going to happen) and then decreases rapidly. I feel that this happened due to human error.

### **Evaluation**

My results were not entirely accurate as when I carried out each experiment and the results didn't turn out the same every time when I repeated it 3 times. I got an anomalous reading for the "edge position of the huddle" as it gained heat at 4 and 6 minutes, this is due to human error. The middle and the penguin stood on its own, also gained heat at, own: 4 minutes and middle at: also 4 minutes. Yet the results that I got did support my prediction.

### **Accuracy**

My results were not entirely accurate as I didn't get the same readings after repeating the investigation 3 times – as I have said before. Also, that I had anomalous readings due to human error. I could have gotten more accurate readings by having 3 people check the readings on the thermometer instead of one person, as the readings could have been different because even 3 seconds could have made the results different. I could have gotten more readings and done the experiment over a longer period of time. I also could have used cotton wool pushed in the top of the test tube as it would have stopped heat escaping as quickly. The results I got were accurate enough to make a firm conclusion. The only pattern, in which I could see, was that the penguin stood on its own, lost the most heat and was the fastest to lose it.

### **Improvements**

I would have changed the method to suit the experiment more, by doing the investigation over a longer period of time as this would make a clearer conclusion with more evidence. I would also get more people to read the results off the thermometer as one person can only read one result at a time and the time difference between reading each one; this could affect the overall conclusion and would be more accurate. I would have decreased the time taken in which I transferred the water from the water bath to the test tubes. And I would have used cotton wool at the top of the test tube so the heat couldn't have escaped.