

**ABUNDANCE, DISTRIBUTION AND SPECIES DIVERSITY OF
INVERTEBRATES COLLECTED BY COMMERCIAL TRAWLER FROM
NEAR SHORE COASTAL AREAS OF KARACHI, PAKISTAN
(NORTHERN ARABIAN SEA).**

**RAOOF MUHAMMAD NIAZI* NAUREEN AZIZ QURESHI **AND MUZAMMIL
AHMED****

***GOVERNMENT CITY COLLEGE,
F. B. AREA, KARACHI.
raoofmniazi@hotmail.com**

****CENTRE OF EXCELLENCE IN MARINE BIOLOGY,
UNIVERSITY OF KARACHI, KARACHI.**

Total number of figures included: 5

Running Title: DIVERSITY OF INVERTEBRATES COLLECTED BY COMMERCIAL TRAWLER

ABSTRACT

The distribution and abundance of benthic macro fauna in near shore waters off Karachi is studied for over a period of one year. For this purpose the marine benthic fauna was collected every month from otter trawl catches of “Hella” fishing trips undertaken by Karachi based commercial shrimp trawlers fishing in depths of about 12 meters. A total of 76 species of invertebrates belonging to 38 families were studied. The distribution and abundance varied with seasons and maximum number of invertebrate species was found during the monsoon season when salinity was low and temperature was high. With increasing salinity and falling temperature the number of invertebrate species in the study area, seemed to decrease. The diversity of invertebrate species collected was studied using the diversity indices and the seasonal variation in the diversity indices was also discussed.

KEY WORDS: ABUNDANCE, DISTRIBUTION, DIVERSITY INDEX, H', COMMERCIAL TRAWLER, COASTAL AREAS OF KARACHI, (NORTHERN ARABIAN SEA).

INTRODUCTION

Commercial trawlers along the coast of Karachi are conducting a lot of bottom trawling. One of the supports of the regional economy and fulfill the needs of protein source of the country. Little information is available in the literature on the bottom dwelling macro-faunas of the coast of Pakistan. There is thus a great gap in our knowledge about our own marine fauna. The main reason for this paucity of information lies in the fact that scientific bottom trawling studies have not been frequently made, and in the occasional cases where undertaken, not much attention has been given to non-commercial or economically unimportant marine organisms. Hence subtidal fauna remains little studied. Compared to the subtidal and benthic marine studies, investigation of the intertidal fauna, flora and their biology are many in number as shown in Azeem-Rahim, (1986). For instance, Ahmed (1977) dealt with the distribution of marine organisms on some rocky, sandy and muddy shores of Karachi region in relation to pollution. Subsequently, Ahmed (1987) dealt with the marine fisheries resources of Indus delta (Northern Arabian Sea).

A research has been conducted in the near shore waters off Karachi, which aims to study the near shore marine benthic fauna, documenting the diversity of the marine fauna. For this purpose the benthic faunas was collected every month for one year. A total of 76 species of invertebrates belonging to 38 families were collected.

MATERIAL AND METHODS

The material was collected by means of an otter trawl operated from a commercial trawler during the period of one year from December 1992 to November 1993. During the study period one trip per month was undertaken. During this period the trawler AL-AHMEDI along the near shore waters off Karachi made a total of 36 hauls. The trawling operations were generally a daylong and are locally known by the name of “Hella” fishery. Random sample from the total catch of the trawl was separated on every trip for subsequent study in laboratory. Identifications of invertebrates were made to species or generic level with the help of available literature. During the monthly observations temperature was measured by the thermometer and salinity by refractometer in the laboratory.

RESULT AND DISCUSSION

On the fishing grounds off Karachi invertebrate fauna was found abundant. A total of 76 species of invertebrates belonging to 38 families were collected. In (Table 1) the month-wise occurrence of three most abundant invertebrate species in the study area were shown among these *Oratosquilla nepa*, *Charybdis callianassa* and *Bulla ampulla* were found to be more numerous than other species in the area, while (Table 2) Shows the dominant number of individuals of invertebrate species in near shore waters off Karachi. Monthly distribution of invertebrate species shown in (Fig.1)

The Shannon and Wiener diversity index was used for the evaluation of species diversity in each collection of the months (Table 3). Diversity index is a mathematical measure of species diversity in a community. Diversity indices provide more information about community composition than simply species richness (i.e., the number of species present). Diversity indices provide important information about rarity and commonness of species in a community. The ability to quantify diversity in this way is an important tool for biologists trying to understand community structure. Diversity commonly depends on the number of species and individuals in the community at a given time and has been mathematically well documented. According to (Fig. 2 & 3). In northeast monsoon season lowest density was observed and higher density in pre monsoon and southwest monsoon seasons (Fig. 4) while in post monsoon season again low density occurred. This pattern shows the diversity of invertebrates high in warmer seasons and low in colder seasons.

Diversity index was calculated by using the Shannon Weiner diversity Index (H) (Shannon Weiner 1963) and equitability or evenness (J) (Pielou 1964) was also calculated. The following formulas were used for the calculations:

$$H = -\sum (P_i) (\log_{10} P_i)$$

$$J = H / H_{\max}$$

Where P_i = Relative percentage of species.

S = Number of species.

H_{\max} = Maximum number of species and is equal to $\log_{10} S$

The study provides valuable information about the occurrence and abundance of marine fauna. It should be continued furthers. The results of the present study provide baseline data for acquiring a better understanding of the macro-benthic life of the near shore waters off Karachi.

Table .1 The month-wise occurrence of three most abundant invertebrate species.

MONTHS	I	II	III
DECEMBER	<i>Thenus orientalis</i>	<i>Sepiella inermis</i>	<i>Calappa lophos</i>
JANUARY	<i>Thenus orientalis</i>	<i>Charybdis callianassa</i>	<i>Calappa lophos</i>
FEBRUARY	<i>Charybdis callianassa</i>	<i>Thenus orientalis</i>	<i>Portunus hastatoides</i>
MARCH	<i>Bulla ampulla</i>	<i>Portunus hastatoides</i>	<i>Stomopneustes variolaris</i>
APRIL	<i>Oratosquilla nepa</i>	<i>Portunus hastatoides</i>	<i>Charybdis callianassa</i>
MAY	<i>Portunus sanguinolentus</i>	<i>Oratosquilla interrupta</i>	<i>Loligo duvauceli</i>
JUNE	<i>Sepiella inermis</i>	<i>Portunus pelagieus</i>	<i>Matuta planipes</i>
JULY	Jellyfish	<i>Loligo duvauceli</i>	<i>Dorippe astuta</i>
AUGUST	<i>Scylla serrata</i>	<i>Renilla reniformis</i>	<i>Babylonia spirata</i>
SEPTEMBER	<i>Portunus sanguinolentus</i>	<i>Natica didyma</i>	<i>Scylla serrata</i>
OCTOBER	<i>Portunus sanguinolentus</i>	<i>Cavernularia obesa</i>	<i>Portunus hastatoides</i>
NOVEMBER	<i>Oratosquilla nepa</i>	<i>Philyra scabriuscula</i>	<i>Oratosquilla hesperia</i>

Table . 2 Dominant invertebrate species in near shore waters off Karachi from December, 1992 to November, 1993. N= Total number of individuals.

S. NO.	SPECIES	N	RANK
1.	<i>Oratosquilla nepa</i>	188	I
2.	<i>Charybdis callianassa</i>	184	II
3.	<i>Bulla ampulla</i>	71	III
4.	<i>Thenus orientalis</i>	51	IV
5.	<i>Portunus sanguinolentus</i>	49	V
6.	<i>Portunus hastatoides</i>	43	VI
7.	<i>Loligo duvauceli</i>	41	VII
8.	<i>Sepiella inermis</i>	40	VIII
9.	<i>Oratosquilla interrupta</i>	35	IX
10.	<i>Calappa lophos</i>	30	X
11.	<i>Charybdis cruciata</i>	22	XI
12.	<i>Astropecten indicus</i>	22	XI




Figure 1. Monthly distributions of invertebrate species in near shore waters of Arabian Sea off Karachi.

Table 3. Diversity and equitability of invertebrate species from near shore waters of Karachi.

MONTHS	DIVERSITY (H')	EQUITABILITY (J)
December	1.20826	1.11961
January	0.9397	0.81989
February	0.72141	0.30903
March	0.79176	0.36946
April	1.0374	0.55326
May	1.21162	0.62965
June	1.27644	0.75130
July	1.25962	0.68288
August	1.00853	0.64803
September	1.18482	0.74467
October	1.06643	0.65697
November	0.82452	0.34095




Figure 2. Monthly Diversity index (H) of Invertebrates from near shore waters of Karachi, (Northern Arabian Sea)




Figure 4. Mean Seasonal Diversity index (H') of invertebrates from near shore Karachi, (Northern Arabian Sea)




Figure 3. Monthly Equitability (J') of Invertebrates from near shore waters of Karachi, (Northern Arabian Sea)

REFERENCES

- AHMED, J. AND KHAN, M. D., 1971. "Pagurids" in the collection of Zoological Survey Department. Rec. Zool. Surv. Pakistan, 2 (2) : 11-16.
- AHMED, M., 1985. Marine Fisheries of Pakistan. Centre of Excellence in Marine Biology. 215 pp.
- AHMED, M. AND MUSTAQUIM, J., 1974. Population structure of four species of Porcellanid crabs (Decapoda: Anomura) occurring on the coast of Karachi. Mar. Biol., 26 : 173-182
- AHMED, *et. al.* , 1972. Crustacean fauna of Pakistan. Marine fauna supplement. Rec. Zool. Surv. Pakistan, 4 (1-2).
- ANONYMOUS, 1972. Marine fauna supplement. Rec. Zool. Surv. Pakistan, 4: 1-68.
- ASHRAF, S. A., 1969. On cephalopods of Pakistan. Rec. Zool. Surv. Pakistan, 1 (2): 1-15.
- ATAUR-RAHIM, M., 1986. Marine science of Northern Arabian Sea. Animal Science Division, Pakistan Agricultural Research Council, Islamabad. 77 pp.
- BIANCHI, G., 1985. Field guide to the commercial marine and brackish water species of Pakistan. FAO Species Identification Sheets for Fishery purposes, Project UNDP/ FAO/ PAK/ 77/ 033, FAO, Rome.

- CHHAPGAR, B. F., 1957. On the marine crabs (Decapoda: Brachyura) of Bombay state. Part I. J. Bombay Nat. Hist. Soc. 54: 399-438.
- CHHAPGAR, B. F., 1957. On the marine crabs (Decapoda: Brachyura) of Bombay state. Part II. J. Bombay Nat. Hist. Soc. 54:503-548.
- CHAUDHRY, A. H., 1924. An account of the Alcyonaria of the Karachi Coast. M. Sc. Thesis, Univ. Punjab.
- FISCHER, W. AND BIANCHI, G. (eds.), 1984. FAO species identification sheets for fishery purpose. Fishing area 51, Western Indian Ocean. Rome. Vols. I-VI.
- HAQ, M. M., 1969. Echinoderms of Pakistan coast. Rec. Zool. Surv. Pakistan, 1 (2): 27-38.
- JEROME, M. E., 1981. A collector's guide to seashells of the world. Published by McGraw-Hill Book Company, New York, 239 pp.
- KHAN, M. A. AND AHMED, M. F., 1975. A check list of Brachyura of Karachi coasts. Rec. Zool. Surv. Pakistan, 7 (1-2).
- KHAN, M. D. AND DASTAGIR, S. G., 1972. On the mollusca: gastropod fauna of Pakistan. Rec. Zool. Surv. Pakistan, II (1): 17-129.
- KHAN, M. D. AND DASTAGIR, S. G., 1979. On the mollusca: pelecypod fauna of Pakistan. Zool. Surv. Dept. Karachi. Printed by the Manager, Printing Corporation of Pakistan Press, Karachi.

MUSTAQUIM, J. AND RABBANI, M., 1976. Species of portunid crab from Karachi. Pakistan J. Sci. Ind. Res., 19: 322-332.

TIRMIZI, N. M., YAQOOB, M. AND SIDDIQUI, F.A., 1982. An illustrated key to the identification of Anomurans (Procellanidae, Albuneidae and Hippidae) of the Northern Arabian Sea. Centre of Excellence in Marine Biology, University of Karachi, Publ. No. 2: 1-29.

