

All four questions on this assignment will be used for assessment purposes.

Question 1 (Unit 9) - 25 marks

Throughout this question, f is an irreducible cubic polynomial in $\mathbf{Q}[t]$.

- (i) Prove that if the complex number γ is a zero of f , then its complex conjugate $\bar{\gamma}$ is also a zero of f . [4]

Let the zeros of f in \mathbf{C} be α, β, γ , and let $L = \mathbf{Q}(\alpha, \beta, \gamma)$. Let $\sigma: L \rightarrow \mathbf{C}$ be defined by $\sigma(x) = \bar{x}$.

- (ii) Prove that $\sigma(L) \subseteq L$ and that $L \subseteq \sigma(L)$. [4]

- (iii) Hence, or otherwise, show that $\sigma: L \rightarrow L$ is a \mathbf{Q} -automorphism of L . [2]

- (iv) Show that if the zeros α, β, γ are *not* all real, then:

(a) $\Gamma(L : \mathbf{Q})$ has a subgroup of order 2; [3]

(b) there is a subfield K of L containing \mathbf{Q} such that $[L : K] = 2$; [3]

(c) $[L : \mathbf{Q}]$ is divisible by 6; [3]

(d) $\mathbf{Q}(\alpha) : \mathbf{Q}$ is not a normal extension. [6]

Question 2 (Unit 10) - 25 marks

The field N is the normal closure of $L : \mathbf{Q}$ in \mathbf{C} , where $L = \mathbf{Q}(\alpha)$ and α is the real fifth root of 7.

- (i) Find $[L : \mathbf{Q}]$ and $|\Gamma(L : \mathbf{Q})|$. [5]

- (ii) Find the number of \mathbf{Q} -monomorphisms from

(a) L into \mathbf{R} ;

(b) L into N ;

(c) L into \mathbf{C} . [4]

- (iii) Prove that $N = L(\beta)$, where $\beta = e^{2\pi i/5}$. [4]

- (iv) Show that $[\mathbf{Q}(\beta) : \mathbf{Q}] = 4$, and hence that $[N : \mathbf{Q}]$ is divisible by 20. [7]

- (v) Show that $[N : \mathbf{Q}(\beta)] \leq 5$, and hence that $[N : \mathbf{Q}] = 20$. [5]

Question 3 (Unit 11) - 25 marks

Let $\varepsilon = e^{2\pi i/8}$. You may assume that the minimum polynomial of ε over \mathbf{Q} is $t^4 + 1$.

- (i) Prove that $\mathbf{Q}(\varepsilon)$ is a splitting field for $t^4 + 1$ over \mathbf{Q} . [3]

Let G be the Galois group $\Gamma(\mathbf{Q}(\varepsilon) : \mathbf{Q})$.

- (ii) Find all the members of G . [5]

- (iii) List all the subgroups of G . [4]

- (iv) How many subfields does $\mathbf{Q}(\varepsilon)$ have? Justify your answer. [5]

- (v) Find all subfields of $\mathbf{Q}(\varepsilon)$, expressing your answers in the form $\mathbf{Q}(x)$, for suitable elements x of $\mathbf{Q}(\varepsilon)$. [8]